

Supersymmetric theories with explicit R -parity violation*

Probir Roy

Tata Institute of Fundamental Research, Mumbai, India

- R_p and motivation for explicit R_p • $MSSM \rightarrow R_p$ MSSM
- Bounds on R_p couplings • Collider-specific phenomenology
- Cosmological implications • Summary and outlook • References

• R_p and motivation for explicit R_p

The superfield spectrum of the Minimal Supersymmetric Standard Model $MSSM$ [1] can be written in usual notation as

$$Q_i = \begin{pmatrix} U_i \\ D_i \end{pmatrix}, \quad L_i = \begin{pmatrix} N_i \\ E_i \end{pmatrix}, \quad H_u = \begin{pmatrix} H_u^+ \\ H_u^0 \end{pmatrix}, \quad H_d = \begin{pmatrix} H_d^0 \\ H_d^- \end{pmatrix}, \quad \bar{U}_i, \bar{D}_i, \bar{E}_i$$

and its superpotential as

$$W_{MSSM} = h_{ij}^u Q_i \cdot H_i U_j + h_{ij}^d Q_i \cdot H_d \bar{D}_j + h_{ij}^e L_i \cdot H_d \bar{E}_j + \mu H_u \cdot H_d. \quad (1)$$

(1) has been constructed so as to conserve R -parity (even for particles, odd for sparticles)

$$R_p \equiv (-1)^{3B+L+2s},$$

B, L and s being baryon no., lepton no. and spin. This is equivalent to the conservation of matter parity which transforms the superfields as

$$(L_i, \bar{E}_i, Q_i, \bar{U}_i, \bar{D}_i) \rightarrow -(L_i, \bar{E}_i, Q_i, \bar{U}_i, \bar{D}_i),$$

$$(H_u, H_d, V_\gamma, V_Z, V_W^+, V_W^-) \rightarrow (H_u, H_d, V_\gamma, V_Z, V_W^+, V_W^-).$$

Three spectacular consequences ensue from R_p -conservation. First, each vertex involving sparticles has them in a pair. Second, the lightest supersymmetric particle (LSP $\tilde{\chi}_1^0$: a weakly interacting neutralino, escaping through detectors, being neutral on cosmological grounds)

*Plenary talk at the Pacific Particle Physics Phenomenology Workshop, Seoul, 31 Oct. – 2 Nov., 1997

is an excellent candidate for cold dark matter. Finally, each sparticle, pair-produced in a collider, would decay within the detector into particles and $\tilde{\chi}_1^0$, the latter being characterized by a hard (E_T) signature.

Recall that R_p -conservation in supersymmetric theories was phenomenologically postulated [2] by the need to avoid catastrophic proton decays induced by R_p vertices. However, such decays need to violate both B and L . So, in place of R_p , it suffices to have another discrete symmetry such as B -parity transforming the superfields as

$$(Q_i, \bar{U}_i, \bar{D}_i) \rightarrow -(Q_i, \bar{U}_i, \bar{D}_i),$$

$$(L_i, \bar{E}_i, H_u, H_d, V_\gamma, V_Z, V_W^+, V_W^-) \rightarrow (L_i, \bar{E}_i, H_u, H_d, V_\gamma, V_Z, V_W^+, V_W^-).$$

Analogously, one can have an L -parity instead. Since our presently scanty knowledge of high scale theories is unable [3] to discriminate between various possible residual discrete symmetries at low energies, it is quite reasonable to consider [4] R_p (but L - or B -conserving) interactions near weak-scale energies. In fact, supersymmetric grand unified theories have been constructed [5] which have R_p interactions present at low energies. There are even stringy scenarios [6] which predict them. On the other hand, a spontaneous violation of R_p requires [7] an extra SM -singlet and hence a nonminimal spectrum; so we will restrict ourselves to explicit R_p -violation. The LSP, in this scenario, is neither stable nor necessarily neutral.

• MSSM $\rightarrow R_p$ MSSM

All renormalizable supersymmetric[†] R_p interactions are incorporated by the addition of new terms in the superpotential

$$W_{MSSM} \rightarrow W_{MSSM} + W_{R_p}, \quad (2a)$$

$$W_{R_p} = \frac{1}{2} \lambda_{[ij]k} L_i \cdot L_j \bar{E}_k + \lambda'_{ijk} L_i \cdot Q_j \bar{D}_k + \frac{1}{2} \lambda''_{i[jk]} \bar{U}_i \bar{D}_j \bar{D}_k + \epsilon_i L_i \cdot H_u, \quad (2b)$$

i, j, k being flavor indices and square brackets on them testifying to antisymmetry in the bracketted indices. The last RHS term in (2) can be “absorbed” in the μ -term of (1) by the transformation $H_d \rightarrow H'_d = (\mu^2 + \epsilon_j \epsilon_j)^{-1/2} (\mu H_d - \epsilon_i L_i)$. This will, in general, lead to $\not\equiv$ terms in the scalar potential and, unless one is careful, can lead to nonzero sneutrino VEVs $\langle \tilde{\nu}_i \rangle \neq 0$. Moreover, the abovementioned “absorption” works at a certain energy scale. At a different energy the μ and ϵ_i parameters will be different, on account of separate renormalization group evolutions, leading to a “reappearance” of the $\epsilon_i L_i \cdot H_u$ terms in W . However, the effects of such terms are smaller [8] than the bounds on the λ -, λ' -, λ'' -couplings that we discuss below and so henceforth we will ignore the $\epsilon_i L_i \cdot H_u$ piece in (2b).

We still have 9 λ -type, 27 λ' -type and 9 λ'' -type, i.e. 45 possible, independent extra, couplings. The R_p Lagrangian density reads

[†]Supersymmetry-breaking R_p terms can occur in the Lagrangian in the form of trilinear scalar products. These have less spectacular effects than the supersymmetric R_p interactions to which we restrict ourselves.

$$\begin{aligned}
\mathcal{L}_{R_p} = & \lambda_{[ij]k} \left[\tilde{\nu}_{iL} \bar{e}_{kR} e_{jL} + \tilde{e}_{jL} \bar{e}_{kR} \nu_{iL} + \tilde{e}_{kR}^* \overline{(\nu_{iL})^C} e_{jL} - \tilde{\nu}_{jL} \bar{e}_{kR} e_{iL} \right. \\
& \left. - \tilde{e}_{iL} \bar{e}_{kR} \nu_{jL} + \tilde{e}_{kR}^* \overline{(\nu_{jL})^C} e_{iL} \right] + \lambda'_{ijk} \left[\tilde{\nu}_{iL} \bar{d}_{kR} d_{jL} + \tilde{d}_{jL} \bar{d}_{kR} \nu_{iL} \right. \\
& \left. + \tilde{d}_{kR}^* \overline{(\nu_{iL})^C} d_{jL} - \tilde{e}_{iL} \bar{d}_{kR} u_{jL} - \tilde{u}_{jL} \bar{d}_{kR} e_{iL} - \tilde{d}_{kR}^* \overline{(e_{iL})^C} u_{jL} \right] + \lambda''_{i[jk]} \epsilon_{\alpha\beta\gamma} \\
& \left[\tilde{u}_{iR\alpha}^* \bar{d}_{kR\beta} d_{jR\gamma}^C + \tilde{d}_{jR\beta} \bar{e}_{kR\gamma} u_{iR\alpha}^C + \tilde{d}_{kR\gamma}^* \overline{(u_{iR\alpha})^C} d_{jR\beta} \right] + h.c.
\end{aligned} \tag{3}$$

Generally, except in a specific case, we take the couplings to be real. Moreover, in analogy with the Yukawa couplings of MSSM, we anticipate a generation hierarchy in their strengths with the third generation ones expected to be the largest. Furthermore, the simultaneous presence of λ or λ' and λ'' type of couplings is extremely unlikely given the lack of observation of either baryon nonconserving nucleon decay or double-nucleon annihilation in nuclei. Indeed, these constraints imply [8]

$$|\lambda'_{11k} \lambda''_{11k}| \lesssim 2 \times 10^{-27} \left(\frac{m_{\tilde{k}}}{100 \text{ GeV}} \right)^2, \tag{4a}$$

$$|\lambda_{233} \lambda''_{112}| \lesssim 10^{-14} \left(\frac{m_{\tilde{k}}}{100 \text{ GeV}} \right)^2, \tag{4b}$$

$$|(\lambda_{\dots} \text{ or } \lambda'_{\dots}) \lambda''_{\dots}| < 10^{-10} \left(\frac{m_{\tilde{k}}}{100 \text{ GeV}} \right)^2, \tag{4c}$$

where $m_{\tilde{k}}$ is the mass of a squark of flavor k (generic in the case of 6c) and $\dots$ stand for any any set of three flavor indices. Because (6) is so stringent, we will henceforth consider either L -violating and B -conserving λ, λ' -couplings or B -violating, L -conserving λ'' -ones.

• Bounds on R_p couplings

In deriving bounds on the absolute values of couplings, we will assume that one R_p operator acts at a time. Known upper bounds [9-14] on various λ - and λ'' type of couplings, along with the physical process which is the source of each bound, are listed in Table 1. Specific references to the original derivation of these bounds may be found in the review by Bhattacharyya [10]. A similar listing has also been done, for whatever bounds [9-15] that exist on the λ' couplings. One remark on the bounds from $D\bar{D}$ mixing and D -decays is in order. In SM no separate information on the unitary matrices $U_L^u, \bar{U}_L^d$, transforming u, d quarks to the mass-diagonal basis, is ever needed. Only the product combination $U_L^{u\dagger} U_L^d \equiv V_{CKM}$ is constrained by experiment. Such is not the case once R_p interactions are included. The bounds above, pertaining to the D -system, have been derived by assuming $U_L^d = 1$.

TABLE 1

Upper bounds on R_p couplings for $\tilde{m} = 100$ GeV. The numbers with $(\star)$ correspond to 2σ limits and those with $(\ddagger)$ are basis-dependent limits.

PART A

Coupling	Bound	Source	Coupling	Bound	Source
$ \lambda_{121} $	0.05 $\star$	CC univ.	$ \lambda''_{112} $	10^{-6}	$NN \rightarrow K$'s
$ \lambda_{122} $	0.05 $\star$	CC univ.	$ \lambda''_{113} $	10^{-5}	$n\bar{n}$ oscillation
$ \lambda_{123} $	0.05 $\star$	CC univ.	$ \lambda''_{123} $	1.25	Pert. unitarity
$ \lambda_{131} $	0.06	$\Gamma(\tau \rightarrow e\nu\bar{\nu})/\Gamma(\tau \rightarrow \mu\nu\bar{\nu})$	$ \lambda''_{212} $	1.25	Pert. unitarity
$ \lambda_{132} $	0.06	$\Gamma(\tau \rightarrow e\nu\bar{\nu})/\Gamma(\tau \rightarrow \mu\nu\bar{\nu})$	$ \lambda''_{213} $	125	Pert. unitarity
$ \lambda_{133} $	0.003	ν_e -mass	$ \lambda''_{223} $	1.25	Pert. unitarity
$ \lambda_{231} $	0.06	$\Gamma(\tau \rightarrow e\nu\bar{\nu})/\Gamma(\tau \rightarrow \mu\nu\bar{\nu})$	$ \lambda''_{312} $	10^{-5}	$n\bar{n}$ oscillation
$ \lambda_{232} $	0.06	$\Gamma(\tau \rightarrow e\nu\bar{\nu})/\Gamma(\tau \rightarrow \mu\nu\bar{\nu})$	$ \lambda''_{313} $	10^{-5}	$n\bar{n}$ oscillation
$ \lambda_{233} $	0.06	$\Gamma(\tau \rightarrow e\nu\bar{\nu})/\Gamma(\tau \rightarrow \mu\nu\bar{\nu})$	$ \lambda''_{323} $	0.50	Γ_ℓ^Z/Γ_h^Z (LEP 1)

PART B

Coupling	Bound	Source	Coupling	Bound	Source	Coupling	Bound	Source
$ \lambda'_{111} $	0.00035	$(\beta\beta)_{0\nu}$	$ \lambda'_{211} $	0.09	R_π	$ \lambda'_{311} $	0.10	$\tau^- \rightarrow \pi^- \nu_\tau$
$ \lambda'_{112} $	0.02 $\star$	CC univ.	$ \lambda'_{212} $	0.09	R_π	$ \lambda'_{312} $	0.10	$\tau^- \rightarrow \pi^- \nu_\tau$
$ \lambda'_{113} $	0.02 $\star$	CC univ.	$ \lambda'_{213} $	0.09	R_π	$ \lambda'_{313} $	0.10	$\tau^- \rightarrow \pi^- \nu_\tau$
$ \lambda'_{121} $	0.035 $\star$	APV	$ \lambda'_{221} $	0.18	D -decay	$ \lambda'_{321} $	0.20 $\ddagger$	$D^0 \bar{D}^0$ mix.
$ \lambda'_{122} $	0.02	ν_e -mass	$ \lambda'_{222} $	0.18	D -decay	$ \lambda'_{322} $	0.20 $\ddagger$	$D^0 \bar{D}^0$ mix.
$ \lambda'_{123} $	0.20 $\ddagger$	$D^0 \leftrightarrow \bar{D}^0$	$ \lambda'_{223} $	0.18	D -decay	$ \lambda'_{323} $	0.20 $\ddagger$	$D^0 \bar{D}^0$ mix.
$ \lambda'_{131} $	0.035 $\star$	APV	$ \lambda'_{231} $	0.22 $\star$	ν_μ d.i.	$ \lambda'_{331} $	0.48	R_τ (LEP)
$ \lambda'_{132} $	0.34	R_e (LEP)	$ \lambda'_{232} $	0.36	R_μ	$ \lambda'_{332} $	0.48	R_τ (LEP)
$ \lambda'_{133} $	0.0007	ν_e -mass	$ \lambda'_{233} $	0.36	R_μ	$ \lambda'_{333} $	0.48	R_τ (LEP)

An interesting recent development has been the derivation [16] of bounds on products of different R_p couplings that are more stringent than the products of the individual bounds available. These follow if, instead of assuming the action of only one operator, we allow the possibility of two different operators acting at two different vertices in a Feynman diagram. Some of these bounds are tabulated below with the corresponding processes cited. Additional results may be found in the recent papers of Jang et al [16].

TABLE 2

Upper bounds on some important product couplings for $\tilde{m} = 100$ GeV.

Combination	Bound	Source	Combination	Bound	Source
$ \lambda'_{11k}\lambda''_{11k} $	10^{-27}	Proton decay	$ \lambda'_{ijk}\lambda''_{lmn} $	10^{-10}	Proton decay
$ \lambda_{1j1}\lambda_{1j2} $	7.10^{-7}	$\mu \rightarrow 3e$	$ \lambda_{231}\lambda_{131} $	7.10^{-7}	$\mu \rightarrow 3e$
$ \text{Im } \lambda'_{i12}\lambda_{i21}^* $	8.10^{-12}	ϵ_K	$ \lambda'_{i12}\lambda'_{i21} $	1.10^{-9}	Δm_K
$ \lambda'_{i13}\lambda'_{i31} $	8.10^{-8}	Δm_B	$ \lambda'_{1k1}\lambda'_{2k2} $	8.10^{-7}	$K_L \rightarrow \mu e$
$ \lambda'_{1k1}\lambda'_{2k1} $	5.10^{-8}	$\mu\text{Ti} \rightarrow e\text{Ti}$	$ \lambda'_{11j}\lambda'_{21j} $	5.10^{-8}	$\mu\text{Ti} \rightarrow e\text{Ti}$
$ \lambda''_{i32}\lambda''_{i21} $	0.008	$\Gamma(B^{\text{ch}} \rightarrow K^{\text{neut.}} K^{\text{ch}})$	$ \lambda''_{i31}\lambda''_{i21} $	0.006	$\Gamma(B^{\text{ch}} \rightarrow K^{\text{neut.}} \pi^{\text{ch}})$

• Collider-specific phenomenology

Signatures for $\tilde{R}_p$ interactions in collider experiments involve two aspects of the concerned processes: first, production of some sparticles and second their specific decay modes. If sparticle pair-production is utilized, as is usually the case, only MSSM vertices are involved in the production process, whereas $\tilde{R}_p$ shows up through the sparticle decays that follow. Single sparticle production is, however, possible with $\tilde{R}_p$ vertices and can be exploited sometimes.

Tevatron $\rightarrow$ *LHC*

In such hadron colliders the possible production and decay-chains are

$$q\bar{q} \text{ or } gg \rightarrow \tilde{q}\tilde{q}^* \rightarrow qq\tilde{\chi}_1^0\tilde{\chi}_1^0,$$

$$\tilde{\chi}_1^0 \rightarrow \ell\bar{\ell}\nu \text{ or } q'\bar{q}\ell.$$

Whether $\tilde{\chi}_1^0$ decays largely into $\ell^+\ell^-\nu$ or two jets and a lepton depends on which of the operators $L_i \cdot L_j \bar{E}_k$ and $L_i \cdot Q_j \bar{D}_k$ in (2b) is dominant. These $\ell\ell\bar{\ell}\bar{\ell} \cancel{E}_T$ or $jjjj\ell\ell$ signals consistute very characteristic signatures [17] of $\tilde{R}_p$. Another process involving $\tilde{R}_p$ interactions in both production and decay is

$$g + d \rightarrow \tilde{u}_j + \ell^-,$$

$$\tilde{u}_j \rightarrow d_k + \bar{\ell},$$

leading to a $\ell\bar{\ell}j$ final state. The lower limits of $O(250)$ GeV, derived from Tevatron data on squark and gluino masses in $MSSM$, get diluted to about $O(100 - 130)$ GeV in R_p $MSSM$. Another important point is that λ' -couplings make additional contributions to top semileptonic decay $t \rightarrow b\ell^+\nu$, leading to a violation of $e - \mu$ universality.

HERA

Resonant squark production in deep inelastic ep scattering is possible via R_p interactions, e.g.

$$\begin{aligned} e^+d &\rightarrow \tilde{u} \rightarrow (e^+d_j, \bar{\nu}u_j, \tilde{\chi}_1^0, u_j, \tilde{\chi}_1^+d_j), \\ e^-u &\rightarrow \tilde{d} \rightarrow (\bar{e}u_j, \bar{\nu}_ed_j, \tilde{\chi}_1^0d_j, \tilde{\chi}_1^-u_j), \\ \tilde{\chi}_1^0 &\rightarrow (\ell^\pm \text{ or } \nu)jj \text{ or } jj\ell, \\ \tilde{\chi}_1^\pm &\rightarrow (\ell^\pm \text{ or } \nu)jj \text{ or } jj\ell, \end{aligned}$$

leading to a variety of final states containing multileptons + $\cancel{E}_T$ or multileptons + jets. For these signals to be observable, assuming that the squark masses are within the energy reach, one will need the relevant λ' couplings to be between [18] $O(10^{-2})$ and $O(10^{-1})$ in magnitude. A point to note is that these are precisely the couplings that are relevant to atomic parity violation (cf. Table 1).

LEP 2 and NLC

In e^+e^- colliders one can look for LSP pair-production followed by R_p decays:

$$\begin{aligned} e^+e^- &\rightarrow \tilde{\chi}_1^0\tilde{\chi}_1^0 \\ \tilde{\chi}_1^0 &\rightarrow \ell^+\ell^-\nu \text{ or } jj\ell. \end{aligned}$$

Thus one can again have quadrilepton + $\cancel{E}_T$ or dilepton + multijet final states. Particularly interesting [14] are final states containing an isolated likesign ditau with multijets. These signals can be visible even if the concerned λ or λ' couplings are as low as $O(10^{-4})$.

• Cosmological implications

Baryogenesis

Any baryon asymmetry, generated at temperatures pertaining to the GUT scale, is in danger of being washed out by sphaleron-induced processes if there are additional $\cancel{B} - \cancel{L}$ interactions at the electroweak scale. The requirement of the preservation of primordial baryon asymmetry was alleged to impose strong ($< 10^{-7}$) constraints on the strengths of R_p interactions which violate $\cancel{B} - \cancel{L}$. However, the conservation of $\frac{1}{3}B - L_i$, where i is any $\cancel{B} - \cancel{L}$ one generation, has been shown [19] to suffice for this purpose – thereby leaving most of the R_p couplings cosmologically unconstrained.

LSP decay

The simplest possible decay mode of the lightest neutralino is $\tilde{\chi}_1^0 \rightarrow \bar{e} u \bar{d}$ with the rate given by Dawson's formula [20] $\tau_{\tilde{\chi}_1^0}^{-1} = (128\pi^2)^{-1} 3\alpha |\lambda'_{121}|^2 M_{\tilde{\chi}_1^0}^5 m_{\tilde{e}}^{-4}$. The requirement of the LSP decaying within a detector can be quantified as $c\gamma_L \tau_{\tilde{\chi}_1^0} < 1m$ and leads to the constraint

$$|\lambda'_{121}| > 1.4 \times 10^{-6} \sqrt{\gamma_L} \left(\frac{m_{\tilde{e}}}{200 \text{ GeV}} \right)^2 \left(\frac{100 \text{ GeV}}{M_{\tilde{\chi}}} \right)^{5/2},$$

where γ_L is the Lorentz boost factor. On the other hand, from the experimental absence of a long-lived relic LSP whose decay would lead to detectable upward going muons, one can infer [21] that $1 \text{ s} < \tau_{\tilde{\chi}} < 10^{17} \text{ yrs}$ leading to the forbidden interval $10^{-10} \not\leq |\lambda, \lambda', \lambda''| \not\leq 10^{-22}$. Note that $\tau_{\tilde{\chi}} > 10^{17} \text{ yrs}$ would make $\tilde{R}_p$ *MSSM* indistinguishable from *MSSM*. Furthermore, the interval $10^{-8} \text{ s} < \tau_{\tilde{\chi}_1} < 1 \text{ s}$ is practically out of observational reach. Of course, with an unstable LSP, one needs another candidate particle (axion?) for cold dark matter.

Summary and outlook

We conclude by emphasizing four main points. (1) There is presently no credible theoretical objection against the presence of $\tilde{R}_p$ interactions near the weak scale. (2) Though many of the $\tilde{R}_p$ couplings have been constrained in Tables 1-3, several (e.g. $\lambda'_{322}, \lambda'_{323}$ etc.) are totally unconstrained. (3) Vigorous seaches are in progress and detection strategies are under formulation for collider signatures of $\tilde{R}_p$ in terms of excess lepton-jet combinations at HERA, violation of $e - \mu$ universality in top semileptonic decay at the Tevatron and isolated ditaus and jets in LEP 2 and NLC. (4) It is highly desirable to find some way of discriminating between different CDM candidates such as a neutralino and an axion.

I compliment the P⁴ workshop organizers for a job well-done.

• References

- [1] H.E. Haber and G.L. Kane, Phys. Rep. **117** (1975) 75.
- [2] G. Farrar and P. Fayet, Phys. Lett. **B76** (1978) 575.
- [3] L.E. Ibanez and G.G. Ross, Phys. Lett. **B260** (1991); Nucl. Phys. **B368** (1992) 3.
T. Banks and M. Dine, Phys. Rev. **D45** (1992) 1424.
- [4] L.J. Hall and M. Suzuki, Nucl. Phys. **B231** (1984) 419.
- [5] D. Brahm and L.J. Hall, Phys. Rev. **D40** (1989) 2449. K. Tamvakis, Phys. Lett. **B383** (1996) 207. G.F. Giudice and R. Rattazzi, hep-ph/9704339.
- [6] M.C. Bento, L.J. Hall and G.G. Ross, Nucl. Phys. **B292** (1987) 400.
- [7] C.S. Aulakh and R.N. Mohapatra, Phys. Lett. **119B** (1982) 136. A. Santamaria and J.W.F. Valle, Phys. Lett. **195B** (1987) 423.

- [8] R. Hempfling, Nucl. Phys. **B478** (1996) 3. B. Mukhopadhyay and S. Roy, Phys. Rev. **D55** (1997) 7020.
- [9] J.L. Goity and M. Sher, Phys. Lett. **B346** (1995) 69. A.Yu. Smirnov and F. Vissani, Phys. Lett. **B380** (1996) 317.
- [10] V. Barger, G.F. Giudice and T. Han, Phys. Rev. **D40** (1989) 2987. G. Bhattacharyya, hep-ph/9709396. H. Dreiner, hep-ph/9707435.
- [11] B. Brahmachari and P. Roy, Phys. Rev. **D50** (1994) 39, err. *ibid.* **D51** (1995) 193.
- [12] D. Chang and W-K. Keung, Phys. Lett. **B389** (1996) 294.
- [13] G. Bhattacharyya, D. Choudhury and K. Sridhar, Phys. Lett. **B355** (1995) 193.
- [14] R.M. Godbole, P. Roy and X. Tata, Nucl. Phys. **B401** (1993) 67.
- [15] M. Hirsch, H.V. Klapdor-Kleingrothaus and S.G. Kovalenko, Phys. Rev. Lett. **75** (1995) 2276. K.S. Babu and R.N. Mohapatra, Phys. Rev. Lett. **75** (1995) 2276.
- [16] D. Choudhury and P. Roy, Phys. Lett. **B378** (1996) 153. J-H. Jang, J.K. Kim and J.S. Lee, Phys. Rev. **D55** (1997) 7296.
- [17] H. Dreiner and G.G. Ross, Nucl. Phys. **B365** (1991) 597. D.P. Roy, Phys. Lett. **B128** (1992) 270.
- [18] D. Choudhury and S. Raychaudhuri, Phys. Lett. **B401** (1997) 54.
- [19] H. Dreiner and G.G. Ross, Nucl. Phys. **B410** (1993) 183.
- [20] S. Dawson, Nucl. Phys. **B261** (1985) 297.
- [21] E.A. Baltz and P. Gondolo, hep-ph/9704411.